


Applied Statistics

Binomial, Poisson, and Gaussian



Troels C. Petersen (NBI)



“Statistics is merely a quantization of common sense”

Probability Density Functions

A Probability Density Function (PDF) $f(x)$ describes the probability of an outcome x :

$$\text{probability to observe } x \text{ in the interval } [x, x+dx] = f(x) dx$$

PDFs are required to be normalized:

$$\int_S f(x) dx = 1$$

The expectation value (aka. mean) and the variance (i.e. standard deviation squared) are then defined as follows:

$$\mu = \int_{-\infty}^{\infty} x f(x) dx$$

$$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$$

Probability Density Functions

The number of PDFs is infinite, and nearly so is the list of known ones:

Discrete distributions [\[edit source | edit beta \]](#)

With finite support [\[edit source | edit beta \]](#)

- The Bernoulli distribution, which takes value 1 with probability p .
- The Rademacher distribution, which takes value ± 1 with equal probability.
- The binomial distribution, which describes the number of successes in a fixed number of independent trials.
- The beta-binomial distribution, which describes the number of successes in a fixed number of trials, where the probability of success is itself a random variable.
- The degenerate distribution at x_0 , where X is certain to take the value x_0 .
- The discrete uniform distribution, where all elements of a finite set are equally likely.
- The hypergeometric distribution, which describes the number of successes in a fixed number of trials, where there is no replacement.
- The Poisson binomial distribution, which describes the number of successes in a fixed number of trials, where the probability of success is not the same for all trials.
- Fisher's noncentral hypergeometric distribution
- Wallenius' noncentral hypergeometric distribution
- Benford's law, which describes the frequency of the digits that appear in the decimal expansion of real numbers.

With infinite support [\[edit source | edit beta \]](#)

- The beta negative binomial distribution
- The Boltzmann distribution, a discrete distribution in statistical mechanics. Special cases include:
 - The Gibbs distribution
 - The Maxwell-Boltzmann distribution
- The Borel distribution
- The extended negative binomial distribution
- The extended hypergeometric distribution
- The **generalized log-series distribution**
- The generalized normal distribution
- The geometric distribution, a discrete distribution with infinite support
- The hypergeometric distribution
- The logarithmic (series) distribution
- The negative binomial distribution or Pascal distribution
- The parabolic fractal distribution
- The Poisson distribution, which describes a very large number of independent trials, each with a small probability of success. The hyper-Poisson, the general Poisson distribution, and the Conway-Maxwell-Poisson distribution, a two-parameter generalization of the Poisson distribution.
- The **Polya-Eggenberger distribution**
- The Skellam distribution, the distribution of the difference of two independent random walks.
- The **skew elliptical distribution**
- The skew normal distribution
- The Yule-Simon distribution
- The zeta distribution has uses in applied statistics
- Zipf's law or the Zipf distribution. A discrete power-law distribution.
- The Zipf-Mandelbrot law is a discrete power-law distribution.

Continuous distributions [\[edit source | edit beta \]](#)

Supported on a bounded interval [\[edit source | edit beta \]](#)

- The Arcsine distribution on $[a, b]$, which is a special case of the beta distribution.
- The Beta distribution on $[0, 1]$, of which the uniform distribution is a special case.
- The Logitnormal distribution on $(0, 1)$.
- The Dirac delta function although not strictly a function, but the notation treats it as if it were a continuous function.
- The continuous uniform distribution on $[a, b]$, where $a < b$.
- The rectangular distribution is a uniform distribution on $[a, b]$.
- The Irwin-Hall distribution is the distribution of the sum of n independent uniform distributions.
- The Kent distribution on the three-dimensional sphere.
- The Kumaraswamy distribution is as versatile as the beta distribution.
- The **logarithmic distribution (continuous)**
- The PERT distribution is a special case of the beta distribution.
- The raised cosine distribution on $[\mu - \pi, \mu + \pi]$
- The reciprocal distribution
- The triangular distribution on $[a, b]$, a special case of the beta distribution.
- The truncated normal distribution on $[a, b]$.
- The U-quadratic distribution on $[a, b]$.
- The von Mises distribution on the circle.
- The von Mises-Fisher distribution on the N -dimensional sphere.
- The Wigner semicircle distribution is important in quantum mechanics.

Supported on semi-infinite intervals, usually $[0, \infty)$

- The Beta prime distribution
- The Birnbaum-Saunders distribution, also known as the Rayleigh distribution
- The chi distribution
 - The noncentral chi distribution
- The chi-squared distribution, which is the sum of independent squared standard normal variables
 - The inverse-chi-squared distribution
 - The noncentral chi-squared distribution
 - The Scaled-inverse-chi-squared distribution
- The Dagum distribution
- The exponential distribution, which describes the time between events in a Poisson process
- The F-distribution, which is the distribution of the ratio of two chi-squared variables which are not necessarily independent
- The noncentral F-distribution
- Fisher's z-distribution
- The folded normal distribution
- The Fréchet distribution
- The Gamma distribution, which describes the time between events in a Poisson process
 - The Erlang distribution, which is a special case of the gamma distribution
 - The inverse-gamma distribution
- The generalized Pareto distribution
- The Gamma/Gompertz distribution
- The Gompertz distribution
- The half-normal distribution

- Hotelling's T-squared distribution
- The inverse Gaussian distribution, also known as the inverse normal distribution
- The Lévy distribution
- The log-Cauchy distribution
- The **log-gamma distribution**
- The log-Laplace distribution
- The log-logistic distribution
- The log-normal distribution, describing variables whose logarithm is normally distributed
- The Mittag-Leffler distribution
- The Nakagami distribution
- The Pareto distribution, or "power law" distribution
- The Pearson Type III distribution
- The **phased bi-exponential distribution** is a special case of the gamma distribution
- The **phased bi-Weibull distribution**
- The Rayleigh distribution
- The Rayleigh mixture distribution
- The Rice distribution
- The shifted Gompertz distribution
- The type-2 Gumbel distribution
- The Weibull distribution or Rosin Rammler distribution, used in grinding, milling and crushing operations.

Supported on the whole real line [\[edit source | edit beta \]](#)

- The Behrens-Fisher distribution, which arises in the analysis of variance
- The Cauchy distribution, an example of a t-distribution with infinite variance
- Chernoff's distribution
- The Exponentially modified Gaussian distribution
- The Fisher-Tippett, extreme value, or log-logistic distribution
 - The Gumbel distribution, a special case of the Fisher-Tippett distribution
- Fisher's z-distribution
- The generalized logistic distribution
- The generalized normal distribution
- The geometric stable distribution
- The Holtsmark distribution, an example of a distribution with a power-law tail
- The hyperbolic distribution
- The hyperbolic secant distribution
- The Johnson SU distribution
- The Landau distribution
- The Laplace distribution
- The Lévy skew alpha-stable distribution or Lévy distribution, Lévy distribution and normal distribution
- The Linnik distribution
- The logistic distribution
- The **map-Airy distribution**
- The normal distribution, also called the Gaussian distribution, the most common distribution
- The Normal-exponential-gamma distribution
- The **Pearson Type IV distribution** (see Pearson Type III distribution)
- The skew normal distribution

- Student's t-distribution, useful for estimating the mean of a normal distribution with unknown variance
 - The noncentral t-distribution
- The type-1 Gumbel distribution
- The Voigt distribution, or Voigt profile, is the convolution of a Gaussian and a Lorentzian distribution
- The Gaussian minus exponential distribution is a special case of the gamma distribution

With variable support [\[edit source | edit beta \]](#)

- The generalized extreme value distribution has three types
- The generalized Pareto distribution has a special case
- The Tukey lambda distribution is either supported on a bounded interval or on the whole real line
- The **Wakely distribution**

Mixed discrete/continuous distributions [\[edit source | edit beta \]](#)

- The rectified Gaussian distribution replaces the normal distribution with a uniform distribution on the negative axis

Joint distributions [\[edit source | edit beta \]](#)

For any set of independent random variables the joint distribution is the product of the individual distributions.

Two or more random variables on the same set [\[edit source | edit beta \]](#)

- The Dirichlet distribution, a generalization of the multinomial distribution
- The Ewens's sampling formula is a probability distribution on the symmetric group
- The Balding-Nichols model
- The multinomial distribution, a generalization of the binomial distribution
- The multivariate normal distribution, a generalization of the normal distribution
- The negative multinomial distribution, a generalization of the negative binomial distribution
- The generalized multivariate log-gamma distribution

Matrix-valued distributions [\[edit source | edit beta \]](#)

- The Wishart distribution
- The inverse-Wishart distribution
- The matrix normal distribution
- The matrix t-distribution

Non-numeric distributions [\[edit source | edit beta \]](#)

- The categorical distribution
- The multinomial distribution

Miscellaneous distributions [\[edit source | edit beta \]](#)

- The Cantor distribution
- The generalized logistic distribution family
- The Pearson distribution family
- The phase-type distribution




And surely more!

Probability Density Functions

An almost complete list of those we will deal with in this course is:

- Gaussian (aka. Normal)
- Poisson
- Binomial (perhaps also Multinomial)
- Students t-distribution
- Uniform
- ChiSquare
- Exponential

You should already know most of these, and the rest will be explained.



Binomial, Poisson, Gaussian

$$f(n; N, p) = \frac{N!}{n!(N-n)!} p^n (1-p)^{N-n}$$

Given **N trials** each with **p chance of success**, how many **successes n** should you expect in total?

This distribution is... **Binomial**:


$$f(n; N, p) = \frac{N!}{n!(N-n)!} p^n (1-p)^{N-n}$$

Mean = Np

Variance = $Np(1-p)$

This means, that the error on a fraction $f = n/N$ is:

$$\sigma(f) = \sqrt{\frac{f(1-f)}{N}}$$



Binomial, Poisson, Gaussian

You count 100 cars, and see 15 red cars. What is your estimate of the fraction (i.e. probability) of red cars and its uncertainty?

- a) 0.150 ± 0.050
- b) 0.150 ± 0.026
- c) 0.150 ± 0.036
- d) 0.125 ± 0.030
- e) 0.150 ± 0.081

From previous page:
$$\sigma(f) = \sqrt{\frac{f(1-f)}{N}}$$

A friend tells you, that 8% of the cars on Blegdamsvej are red. What is the chance of that? Could he be right?

Binomial, Poisson, Gaussian

You count 100 cars, and see 15 red cars. What is your estimate of the fraction (i.e. probability) of red cars and its uncertainty?

- a) 0.150 ± 0.050
- b) 0.150 ± 0.026
- c) 0.150 ± 0.036
- d) 0.125 ± 0.030
- e) 0.150 ± 0.081

$$(0.150 - 0.080) / 0.036 = 1.9 \sigma$$



From previous page:
$$\sigma(f) = \sqrt{\frac{f(1-f)}{N}}$$

A friend tells you, that 8% of the cars on Blegdamsvej are red. What is the chance of that? Could he be right?

Binomial, Poisson, Gaussian

Requirements to be Binomial:

- Fixed number of trials, N
- Independent trials.
- Only two outcomes (success / failure).
- Constant probability of success / failure.

If number of possible outcomes is more than two \Rightarrow **Multinomial distribution.**

Examples of Binomial experiments:

- Tossing a coin 20 times counting number of tails.
- Asking 200 people if they watch sports on TV.
- Rolling a die to see if a 6 appear (Multinomial for all outcomes!).
- Asking 100 die-hards from Enhedslisten, if they would vote for Konservative at the next election!

Examples which aren't Binomial experiments:

- Rolling a die until a 6 appears (not fixed number of trials).
- Drawing 5 cards for a poker hand (no replacement \Rightarrow not independent)

Binomial, Poisson, Gaussian

If $N \rightarrow \infty$ and $p \rightarrow 0$, but $Np \rightarrow \lambda$ then a Binomial approaches a Poisson:

$$f(n; \nu) = \frac{\nu^n}{n!} e^{-\nu}$$

In reality, the approximation is already quite good at e.g. $N=50$ and $p=0.1$.


The Poisson distribution only has one parameter, namely λ .

Mean = λ

Variance = λ

So the error on a number is...

...the square root of that number!



Binomial, Poisson, Gaussian

If $N \rightarrow \infty$ and $p \rightarrow 0$, but $Np \rightarrow \lambda$ then a Binomial approaches a Poisson:

$$f(n, \nu) = \frac{\nu^n}{n!} e^{-\nu}$$

In reality, the approximation is already quite good at e.g.

$N=50$ and $p=0.1$.


The Poisson distribution only has one parameter, namely λ .

Mean = λ

Variance = λ

So the error on a number is...

The error on a (Poisson) number... is the square root of that number!!!



...the square root of that number!

The error on a
(Poisson) number...
is the square root
of that number!!!

Note: The sum of two Poissons with λ_a and λ_b is a new Poisson with $\lambda = \lambda_a + \lambda_b$.
(See Barlow pages 33-34 for proof)

Binomial, Poisson, Gaussian


Last year Denmark had 180 deaths in traffic. What is roughly the uncertainty on this number?

- a) 5
- b) 9
- c) 14
- d) 25
- e) Cannot be determined from the information given.

Binomial, Poisson, Gaussian

If $\lambda \rightarrow \infty$ and many other cases...

...and for λ , 20 is enough!




Binomial, Poisson, Gaussian

If $\lambda \rightarrow \infty$ and many other cases...

...and for λ , 20 is enough!

Poisson and Gaussian distribution comparison




Binomial, Poisson, Gaussian

If $\lambda \rightarrow \infty$ and many other cases...


...and for λ , 20 is enough!

Poisson and Gaussian distribution comparison



Binomial, Poisson, Gaussian

"If the Greeks had known it, they would have deified it."



"If the Greeks had known it, they would have deified it. It reigns with serenity and in complete self-effacement amidst the wildest confusion. The more huge the mob and the greater the apparent anarchy, the more perfect is its sway. It is the supreme Law of Unreason. Whenever a large sample of chaotic elements are taken in hand and marshaled in the order of their magnitude, an unsuspected and most beautiful form of regularity proves to be latent all along." [Karl Pearson]

Binomial, Poisson, Gaussian

The Gaussian defines the way we consider uncertainties.

| Range | Inside | Outside |
|---------------|---------------|-----------|
| $\pm 1\sigma$ | 68 % | 32 % |
| $\pm 2\sigma$ | 95 % | 5 % |
| $\pm 3\sigma$ | 99.7 % | 0.3 % |
| $\pm 5\sigma$ | 99.99995 % | 0.00005 % |

